



Citation: Surachest Aiumsumang, Patcharaporn Chaiyasan, Kan Khoomsab, Weerayuth Supiwong, Alongklod Tanomtong Sumalee Phimphan (2022) Comparative chromosome mapping of repetitive DNA in four minnow fishes (Cyprinidae, Cypriniformes). *Caryologia* 75(2): 71-80. doi: 10.36253/caryologia-1523

Received: December 02, 2021

Accepted: May 20, 2022

Published: September 21, 2022

Copyright: © 2022 Surachest Aiumsumang, Patcharaporn Chaiyasan, Kan Khoomsab, Weerayuth Supiwong, Alongklod Tanomtong Sumalee Phimphan. This is an open access, peer-reviewed article published by Firenze University Press (<http://www.fupress.com/caryologia>) and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Competing Interests: The Author(s) declare(s) no conflict of interest.

Comparative chromosome mapping of repetitive DNA in four minnow fishes (Cyprinidae, Cypriniformes)

SURACHEST AIUMSUMANG¹, PATCHARAPORN CHAIYASAN², KAN KHOOMSAB³, WEERAYUTH SUPIWONG⁴, ALONGKLOD TANOMTONG² SUMALEE PHIMPHAN^{1,*}

¹ Biology Program, Faculty of Science and Technology, Phetchabun Rajabhat University, Phetchabun 67000, Thailand

² Program of Biology, Faculty of Science, Khon Kaen University, Muang, Khon Kaen 40002, Thailand

³ Education Science Program, Faculty of Science and Technology, Phetchabun Rajabhat University, Phetchabun 67000, Thailand

⁴ Applied Science Program, Faculty of Interdisciplinary Studies, Khon Kaen University, Nong Khai Campus, Muang, Nong Khai 43000, Thailand

*Corresponding author. E-mail: sumalee.phi@pcru.ac.th

Abstract. The present study focused on the repetitive DNA of the chromosome in four minnow fishes from the genera *Danio* Hamilton, 1822, *Devario* Heckel, 1843 and *Rasbora* Bleeker, 1859. Chromosomes were analysed using fluorescence *in situ* hybridization (FISH) with microsatellite probes including (CA)₁₅, (CAC)₁₀, (CGG)₁₀, (GC)₁₅ and (TA)₁₅ staining. All species retained the diploid chromosome number 2n = 50 in male and female. The microsatellite sequences were mapped in the chromosomes of *Danio albolineatus* (Blyth, 1860), *Devario regina* (Fowler, 1934), *Rasbora aurotaenia* Tirant, 1885 and *R. paviana* Tirant, 1885. In most cases, the microsatellite was dispersed in the chromosome with conspicuous markings in the telomeric region and the whole genome, which suggests that sequences contribute to chromosome structure and may have played a role in the relationship of this fish group. The comparative genome mapping data presented here provide novel information on the structure and organisation of the repetitive DNA region of the minnow's genome and contribute to a better understanding of the genomes of these minnows.

Keywords: Cyprinidae, Danioninae, FISH, microsatellite.

INTRODUCTION

The level of molecular cytogenetics plays an important role in precise characterisation of the structure of fish genomes (Cioffi and Bertollo 2012). The family Cyprinidae is the most abundant and globally widespread family of freshwater fish, comprising 3,000 species are one of the largest groups (Eschmeyer and Fong 2015). Thailand is one of the important areas for fish fauna in terms of both diversity and endemism. The total number of fish spe-

cies in Thai waters is over 2,700 with about 2,000 marine and 720 freshwater species (Vidthayanon 2005). *Danio albolineatus* (Blyth, 1860), *Devario regina* (Fowler, 1934), *Rasbora aurotaenia* Tirant, 1885 and *R. paviana* Tirant, 1885 are four of the species of minnows, belonging to the family Cyprinidae (subfamily Danioninae-Danionini). They are tropical freshwater fish of minor commercial importance, which are native in Thailand. Their distributions include the Mekong, Chao Phraya, and Meklong Basins (Froese and Pauly 2012) and they can be easily found in large and small rivers, ponds, ditches, lakes, paddy field, and swamps. It rarely occurs in low oxygen waters (Brittan 1954; 1971; 1998). They could be used to assess if they were sensitive to change in environmental problems and aquatic pollution (Blazer 2002; Frame and Dickerson 2006; Raskovic et al. 2010; Yenchum 2010; Reddy and Rawat 2013). However, cytogenetic studies in minnows are quite scarce, in which only conventional technique reported to determine chromosome number and karyotype composition has been performed.

Up to the present time, cytogenetic studies on subfamily Danioninae (Cyprinidae, Osteichthyes) with 20 genera and about 335 species have been undertaken for only 21 species from three genera (*Danio*, *Devario* and *Rasbora*) as yet, only conventional cytogenetics have been applied to determine chromosome numbers and karyotype complements. Diploid chromosomes number (2n) varies between 48-50. Cytogenetic studies of the genera *Danio*, *Devario* and *Rasbora* have been reported in Table 1. From the previous reports, in most cases, more descriptions of karyotypes seem to be inconclusive when they are not used in combination with other methods to produce more accurate chromosome markers. More recently, molecular cytogenetics begun to be implemented in a finer-scale characterization of karyotype structures in certain Cyprinid taxa (Spoz et al. 2014; Saenjundaeng et al. 2018; Phimphan et al. 2020). More specifically, FISH-based repetitive DNA mapping, multiple DNA copies of repetitive DNAs are a large substantial portion of the genome of eukaryotes that can be generally classified into two main classes: tandem repeats, such as the multigene families and satellite DNAs; also, there are dispersed elements, such as transposons and retrotransposons, known as transposable elements (TEs) (Jurka et al. 2005). Repetitive DNA sequences display a high degree of polymorphism because of the variation in the number of repetitive units, which results from specific evolutionary dynamics. The taxonomic status by DNA genome as well as the phylogenetic relationships of Danioninae species is confirmed and well established based on previous study (Rüber et al., 2007; Tang et al., 2010).

Recently, molecular cytogenetic studies, using fluorescence *in situ* hybridization (FISH) for mapping of repetitive DNA sequences, have provided important contributions to the characterisation of biodiversity and evolution of divergent fish groups (Cioffi and Bertollo 2012). Moreover, some microsatellite repeats are species-specific characters amongst some fish groups (Cioffi et al. 2015). An important role of repetitive DNAs in genome evolution has been reported for different fish groups (Cioffi and Bertollo 2012; Cioffi et al. 2010, 2015; Moraes et al. 2017; 2019; Sassi et al. 2019; Terencio et al. 2013; Yano et al. 2014).

Thus, the present study is the report on chromosomal characteristics of FISH mapping of the microsatellite repeats in *Da. albolineatus*, *De. regina*, *R. aurotaenia* and *R. paviana* by using molecular cytogenetic protocols. The knowledge gained can provide cytogenetic information potentially useful for further study in this family.

MATERIAL AND METHODS

Individuals from both sexes of the four minnows were collected for analyses from river basins in, Thailand (Table 2 and Fig. 1). The fishes were transferred to laboratory aquaria and kept under standard conditions for three days before the experiments. The procedures followed ethical protocols, as approved by the Ethics of Animal Experimentation of the National Research Council of Thailand U1-04498-2559.

Preparation of fish chromosomes was from kidney cells (Phimphan et al. 2020). The chromosomes were stained with Giemsa's Solution for 10 min. Metaphase figures were analysed according to the chromosome classification of Levan et al. (1964). Chromosomes were classified as metacentric (m), submetacentric (sm), subtelocentric (st) or acrocentric (a). The Fundamental number, NF (number of chromosome arms) is obtained by assigning a value of two (2) to metacentric and submetacentric chromosomes and one (1) to subtelocentric and acrocentric chromosomes.


FISH was performed under stringent conditions on metaphase chromosome spreads with microsatellite (CA)₁₅, (CAC)₁₀, (CGG)₁₀, (GC)₁₅ and (TA)₁₅ probes (Kubat et al. 2008; Liehr 2009) which were directly labelled with Cy3 at 5' terminal during synthesis by Sigma (St. Louis MO, USA). FISH, under stringent conditions on mitotic chromosome spreads (Pinkel et al. 1986), was carried out by previous protocols as reported by Supiwong et al. (2019) and Yano et al. (2017). The hybridization signals were checked and analysed on an

Table 1. Reviews of cytogenetic reports in the genera *Danio*, *Devario* and *Rasbora*. (2n = diploid number, m = metacentric, sm = submetacentric, st = subtelocentric, a = acrocentric, NORs = nucleolar organizer regions, NF = fundamental number, + = positive - = not available).

Genus/Species	2n	NF	Karyotype formula	FISH	Reference
<i>Danio rerio</i>	48	-	-	-	Post (1965)
	50	100	50m	-	Endo and Ingalls (1968)
	50	100	10m + 40sm	-	Fontana et al. (1970)
	50	100	10m + 12sm + 28a	-	Rishi (1976)
	50	100	16m + 32sm + 2a	-	Schreeb et al. (1993)
	50	100	12m + 26sm + 12a	-	Pijnacker and Ferwerda (1995)
	50	100	4m + 16sm + 30a	-	Daga et al. (1996)
	50	100	4m + 16sm + 30a	-	Gornung et al. (1997)
	50	100	F: 7m + 7sm + 36a M: 6m + 8sm + 36a	-	Sharma et al. (1998)
	50	100	12m + 26sm + 12a	-	Ueda and Naoi (1999)
	50	100	4m + 30sm + 16a or*	-	Amores and Postlethwait (1999)
	50	100	4m + 20sm + 26a	-	
	50	100	4m + 16sm + 30a	+	Phillips and Reed (2000)
	50	100	4m + 16sm + 30a	+	Sola and Gornung (2001)
	<i>Da. Roseus</i>	50	100	8m+34sm+8a	-
<i>Da. albolineatus</i>	50	99	10m+39sm+1a	-	Arai (2011)
	50	100	8m + 14sm + 28a	+	Present study
<i>Devario laoensis</i>	50	100	6m+10sm+34a	-	Aiumsumang et al. (2021) Kaewtip et al. (2021)
	50	96	6m+20sm+20a+4t	-	
<i>De. aequipinnatus</i>	50	96	6m+34sm+6st+4t	-	Sukham et al. (2013)
<i>De. regina</i>	50	100	6m+12sm+32a	-	Aiumsumang et al. (2021)
	50	100	6m+12sm+32a	+	Present study
<i>Rasbora agilis</i>	50	100	24m+26sm	-	Donsakul et al. (2009)
<i>R. aurotaenia</i>	50	92	14m+26sm+2a+8t	-	Seetapan and Moeikum (2004)
	50	98	8m+16sm+24a+2t	-	Aiumsumang et al. (2012)
	50	98	8m+16sm+24a+2t	+	Present study
<i>R. borapetensis</i>	50	88	24m+14sm+12t	-	Donsakul et al. (2005)
<i>R. buchani</i>	50	100	30m+18sm+2a	-	Manna and Khuda-Bukhsh (1977)
<i>R. caudimaculata</i>	50	98	20m+26sm+2a+2t	-	Donsakul and Magtoon (2002)
<i>R. daniconius</i>	50	80	18m+6sm+6a+20t	-	Khuda-Bukhsh et al. (1979)
	50	92	32m+8sm+2a+8t	-	Donsakul et al. (2005)
<i>R. dorsiocellata</i>	50	92	18m+24sm+8t	-	Donsakul et al. (2009)
<i>R. einthovenii</i>	50	94	6m+30sm+8a+6t	-	Donsakul et al. (2005)
	50	100	16m+18sm+16a	-	Yeesaem et al. (2019)
<i>R. heteromorpha</i>	48	-	-	-	Post (1965)
	48	74	14m+10sm+2a+22t	-	Donsakul et al. (2005)
<i>R. myersi</i>	50	90	20m+14sm+6a+10t	-	Donsakul and Magtoon (2002)
<i>R. paviei</i>	50	100	10m+24sm+16a	-	Donsakul and Magtoon (2002)
<i>R. paviana</i>	50	98	8m+16sm+24a+2t	-	Aiumsumang et al. (2021)
	50	98	8m+16sm+24a+2t	+	Present study
<i>R. retrodorsalis</i>	50	88	26m+10sm+2a+12t	-	Donsakul and Magtoon (2002)
<i>R. rubrodorsalis</i>	50	82	16m+16sm+18t	-	Donsakul et al. (2009)
<i>R. sumatrana</i>	50	94	26m+16sm+2a+6t	-	Donsakul and Magtoon (1995)
<i>R. trilineata</i>	48	-	-	-	Post (1965)
	50	94	26m+16sm+2a+6t	-	Donsakul et al. (2005)

Table 2. Collection sites of the analyzed species show the sample number.

Species	Number of specimens in site sampling					
	Mae Khong Basin	Sirindhorn Peat Swamp Forest	Pasak Basin	Chi Basin	Chao Phaya Basin	Songkhram Basin
<i>Danio albolineatus</i>	-	-	10 ♀ 10 ♂	04 ♀ 05 ♂	05 ♀ 05 ♂	-
<i>Devario regina</i>	05 ♀ 06 ♂	06 ♀ 08 ♂	-	-	-	-
<i>Rasbora aurotaenia</i>	-	-	-	-	08 ♀ 07 ♂	05 ♀ 08 ♂
<i>Rasbora paviana</i>	05 ♀ 08 ♂	03 ♀ 04 ♂	05 ♀ 07 ♂	04 ♀ 05 ♂	-	-

**Figure 1.** Map showing the collection sites of *Danio albolineatus* [red circles], *Devario regina* [blue circles], *Rasbora aurotaenia* [green circles] and *Rasbora paviana* [yellow circles] for studied herein.

epifluorescence microscope Olympus BX50 (Olympus Corporation, Ishikawa, Japan).

RESULTS

Diploid number and fundamental number of Da. albolineatus, De. regina, R. aurotaenia and R. paviana

The four minnow fishes have the same diploid number of $2n = 50$. Although the minnow fishes analysed

share the same $2n$, there are differences in the fundamental number (NF) i.e. *Da. albolineatus* and *De. regina* NF = 100, while for the two *Rasbora*, NF = 98. The karyotype complements of *D. albolineatus* composed of $8m + 14sm + 28a$, *D. regina* was $m6+sm10+a34$, while *R. aurotaenia* and *R. paviana* were $8m+16sm+24a+2t$ (Figs. 2A-D).

Patterns of the microsatellite repeat in the genomes of Da. albolineatus, De. regina, R. aurotaenia and R. paviana

The result of the mapping of the microsatellite repeats $(CA)_{15}$ show that hybridization signals are abundantly distributed on telomeric regions in all species (Figs. 3E-H), $(CAC)_{10}$ showing moderate abundance in *Da. albolineatus* and *R. paviana*, while *De. regina* and *R. aurotaenia* were not detected (Figs. 3I-L). The region hybridized $(CGG)_{10}$ of *Da. albolineatus* and *R. paviana* identified a partial genome, *De. regina* has a hybridization pattern throughout the genomes and *R. paviana* was not detected (Figs. 3M-P). $(GC)_{15}$ hybridized in *Da. albolineatus* and *R. paviana* as whole genomes, *Da. albolineatus* as telomeric regions and *R. paviana* was not detected (Figs. 3Q-T). The microsatellite $(TA)_{15}$ probe showed hybridization on the whole genomes of *Da. albolineatus* and *De. regina*, while two *Rasbora* were not detected (Figs. 3U-X) (Table 3).

DISCUSSION

Diploid number of and fundamental number Da. albolineatus, De. regina, R. aurotaenia and R. paviana

Da. albolineatus had $2n = 50$ which is in accordance with one single previous report (Arai 2011) and the same in other species in genus *Danio* (Post 1965; Endo and Ingalls 1968; Fontana et al. 1970; Rishi 1976; Schreeb et al. 1993; Pijnacker and Ferwerda 1995; Daga et al. 1996; Gornung et al. 1997; Sharma et al. 1998; Ueda and Naoi 1999; Amores and Postlethwait 1999; Phillips




Figure 2. Karyotype of *Danio albolineatus* [A], *Devario regina* [B], *Rasbora aurotaenia* [C], *Rasbora paviana* [D], m = metacentric, sm = submetacentric, a = acrocentric and t = telocentric chromosomes.

Table 3. Molecular cytogenetic studies on four minnow fishes. [2n = diploid chromosome number, NF fundamental number (number of chromosome arm)].

Species	2n	NF	(CA) ₁₅	(CAC) ₁₀	(CGG) ₁₀	(GC) ₁₅	(TA) ₁₅
<i>Da. albolineatus</i>	50	100	telomere	partial genome	partial genome	whole genome	whole genome
<i>De. regina</i>	50	100	telomere	not detected	whole genome	telomere	whole genome
<i>R. aurotaenia</i>	50	98	telomere	not detected	not detected	partial genome	not detected
<i>R. paviana</i>	50	98	telomere	partial genome	partial genome	whole genome	not detected

and Reed 2000; Sola and Gornung 2001), *Da. Roseus*: 2n=50 (Kaewtip et al. 2021) and *Da. albolineatus*: 2n=50 (Arai 2011). The 2n of *De. regina* is the same as that of *De. laoensis* Pellegrin & Fang, 1940 (Aiumsumang et al. 2021; Kaewtip et al. 2021) and *De. aequipinnatus* reported by Sukham et al. (2013). *R. aurotaenia* has 2n=50 according to Seetapan and Moeikum (2004) and Aiumsumang et al. (2012). The diploid chromosome number of *R. paviana* was 50, which is the same as that from the previous study by Aiumsumang et al. (2021). This 2n is considered the same as for the other species of subfamily Danioninae (*Danio*, *Devario* and *Rasbora*).

Patterns of microsatellite repeats in the genomes of *Da. albolineatus*, *De. regina*, *R. aurotaenia* and *R. paviana*

The chromosomal distribution of repetitive DNA elements revealed remarkable differences amongst the

analysed species. In this study, the number and distribution of microsatellite sequence (CA)₁₅, (CAC)₁₀, (CGG)₁₀, (GC)₁₅ and (TA)₁₅ were not conserved among four analyzed species. The microsatellite sequence (CA)₁₅ was mapped on chromosomes of *Da. albolineatus*, *De. regina*, *R. aurotaenia* and *R. paviana*, and abundantly located and distributed in all chromosomes, usually in telomeric regions and a few chromosome pairs showed the strongest signal intensities, these not really being suited to serve as chromosomal markers. Microsatellites are usually located in the heterochromatic regions (telomeres/centromeres) of fish genomes, where a significant fraction of repetitive DNA is localized (Cioffi and Bertollo, 2012). Indeed, this distribution pattern is found in *Epalzeorhynchus frenatum* (Fowler, 1934), *Puntigrus partipentazona* (Fowler, 1934), *Scaphognathops bandanensis* Boonyaratpalin & Srirungroj, 1971 (Phimphan et al, 2020); *Catlocarpio siamensis* Boulenger, 1898 and *Probarbus jullieni* Sauvage, 1880 (Saenjundaeng et




Figure 3. FISH using DAPI (A-D), $[CA]_{15}$ (E-H), $[CAC]_{10}$ (I-L), $[CGG]_{10}$ (M-P), $[GC]_{15}$ (Q-T), $[TA]_{15}$ (U-Z) of *Danio albolineatus*, *Devario regina*, *Rasbora aurotaenia* and *Rasbora paviana*, respectively. Scale bar 5 μ m.

al. 2018); *Clarias* species (Maneechot et al. 2016) and *Channa micropeltes* (Cuvier, 1831) (Cioffi et al. 2015). This distribution pattern are differences known for *Channa gachua* (Hamilton, 1822), *C. lucius* (Cuvier, 1831), *C. striata* (Bloch, 1793) (Cioffi et al. 2015), *Hemibagrus wyckii* (Bleeker, 1858) (Supiwong et al. 2017) and *Monopterus albus* (Zuiew, 1793) (Supiwong et al. 2019). For the mapping of (CAC)₁₀ repeats in *Da. albolineatus* and *R. paviana*, partial genomes were identified, while *De. regina* and *R. aurotaenia* were not detected. In *Da. albolineatus*, *De. regina*, *R. aurotaenia* and *R. paviana*, most microsatellites (CGG)₁₀ and (CG)₁₅ were abundantly distributed in all chromosomes with the exception of (CGG)₁₀ in *R. aurotaenia*, this not being a signal probe. Nevertheless, comparative analyses among the species indicate that the microsatellites have also preferential zones of accumulation at telomeric heterochromatin. For *De. regina* the accumulation of microsatellites (GC)₁₅ in all chromosomal pair could point out a species-specific type of heterochromatin is similar to previous reports in *Channa micropeltes* (Cioffi et al. 2015). In *Da. albolineatus* *R. aurotaenia* and *R. paviana* the same microsatellites are not only located at the telomeric region of some chromosomes, but they are also found at the centromeres of the chromosome pair. (TA)₁₅ repeats in *Da. albolineatus* and *De. Regina*, displaying high accumulations at the whole genome, while not detected in the *Rasbora* group.

Repetitive DNA sequences display a high degree of polymorphism because of the variation in the number of repetitive units, which results from a specific evolutionary dynamic. Amongst these elements, microsatellites (or short tandem repeats) are the most polymorphic and consist of short sequences of one to six nucleotides repeated in tandem throughout the DNA (Tautz and Renz 1984). Due to their supposed neutral evolution, these molecular markers have been widely used in population genetics, to identify taxonomic limits and in hybridization and forensic studies (Goldstein and Pollock 1997; Filcek et al. 2005; Racey et al. 2007; McCusker et al. 2008) and can be used to spot genomic evolution as previously been reported for different fish groups (Cioffi et al. 2010; Cioffi and Bertollo 2012; Terencio et al. 2013; Yano et al. 2014; Cioffi et al. 2015; Moraes et al. 2017; 2019; Sassi et al. 2019).

The comparative study on four species showed that the diploid chromosome is the same, but the patterns of microsatellite repeat on chromosomes have differences amongst them. Thus, the molecular cytogenetic data may be a tool for classification of fish species where there is similar morphology, such as the stripe. Overall, it is believed that the microsatellites have specific zones of accumulation in genomes, preferentially in hetero-

chromatic regions (Supiwong et al. 2014). In fact, microsatellites are located in the heterochromatic regions (telomeres, centromeres and in the sex chromosomes) of fish genomes (Cioffi and Bertollo 2012, including the present study). However, the distribution of microsatellites was not only restricted to heterochromatin, but also dispersed in euchromatic regions of the chromosomes (Getlekha et al. 2016). Additionally, such variability is also reinforced by the dynamism of repetitive elements in the genome, especially by the differential distribution and accumulation of rDNA sequences amongst chromosomes (Cioffi et al. 2015). Although not yet completely understood, this marked diversity is likely linked to the lifestyle of these fishes and to population fragmentation, as already identified for other fish species.

CONCLUSIONS

Our study is the first one to offer reliable chromosomal data for *Da. albolineatus*, *De. regina*, *R. aurotaenia* and *R. paviana* by molecular cytogenetic protocols, this study were useful tools in highlighting the remarkable chromosomal diversification which was characterised in the four minnow fishes. Besides, data from comparative genomic hybridization experiments also highlighted an advanced stage of repetitive DNA divergence, evidencing their evolutionary diversification.

ACKNOWLEDGMENTS

This study was supported by the National Research Council of Thailand under the Phetchabun Rajabhat University (Grant No. FRB640052/04) and the Post-Doctoral Training Program from Research Affairs and Graduate School, Khon Kaen University, Thailand (Grant no. 59255).

REFERENCES

- Aiumsumang S, Phimphan S, Suwannapoom C, Chaiyasan P, Supiwong W, Tanomtong A. 2021. A comparative chromosome study on five Minnow fishes (Cyprinidae, Cypriniformes) in Thailand. *Caryologia* 74(1): 87–94. <https://doi.org/10.36253/caryologia-1017>
- Amores A, Postlethwait JH. 1999. Banded chromosomes and the zebrafish karyotype. *Methods in Cell Biology* 60: 323–338. [https://doi.org/10.1016/s0091-679x\(08\)61908-1](https://doi.org/10.1016/s0091-679x(08)61908-1)

- Arai R. 2011. Fish Karyotypes: A Check List. Springer Japan, Tokyo. <https://doi.org/10.1007/978-4-431-53877-6>
- Blazer VS. 2002. Histopathological assessment of gonadal tissue in wild fishes. *Physiology Biochemistry* 26: 85–101. <https://doi.org/10.1023/A:1023332216713>
- Brittan MR. 1954. A revision of the Indo-Malayan fresh-water fish genus *Rasbora*. Institute of Science and Technology. Manila Monogr 3: 1–224.
- Brittan MR. 1971. *Rasbora*: A Revision of the Indo-Malayan Fresh-Water Fish Genus *Rasbora*. T.F.H. Publications, Neptune City.
- Brittan MR. 1998. *Rasboras: Keeping and Breeding Them in Captivity*. T.F.H. Publications, Neptune City.
- Cioffi MB, Bertollo LAC. 2012. Distribution and evolution of repetitive DNAs in fish. In: Garrido-Ramos, M. A. (ed.) *Repetitive DNA*, Vol. 7. Karger, Basel, pp. 197–221. <https://doi.org/10.1159/000337952>
- Cioffi MB, Bertollo LAC, Villa MA, Oliveira EA, Tanomtong A, Yano CF, Supiwong W, Chaveerach A. 2015. Genomic organization of repetitive DNA elements and its implications for the chromosomal evolution of channid fishes (Actinopterygii, Perciformes). *PLoS One*. 10(6): e0130199. <https://doi.org/10.1371/journal.pone.0130199>
- Cioffi MB, Martins C, Bertollo LAC. 2010. Chromosome spreading of associated transposable elements and ribosomal DNA in the fish *Erythrinus erythrinus*. Implications for genome change and karyoevolution in fish. *BMC Evolution. Biology* 10(1): 271–279. <https://doi.org/10.1186/1471-2148-10-271>.
- Daga RR, Thode G, Amores A. 1996. Chromosome complement, C banding, Ag-NOR and replication banding in the zebrafish *Danio rerio*. *Chromosome Research* 4(1): 29–32. <https://doi.org/10.1007/BF02254941>
- Donsakul T, Rangsiruji A, Magtoon W. 2009. Karyotypes of five cyprinid fishes (Cyprinidae, Danioninae-Danionini): *Rasbora agilis*, *R. dorsicellata*, *R. rubrodorsalis*, *Boraras maculate* and *B. urophthalmoides* from Thailand. In Proceedings of the 47th Kasetsart University Annual Conference, Kasetsart. Bangkok, Thailand. 320–327.
- Donsakul T, Magtoon W, Rangsiruji W. 2005. Karyotype of five species of fish (Pla siew) in subfamily Rasboranae. 31st Congress on Science and Technology of Thailand at Suranaree University of Technology. Nakhon Ratchasima Province Thailand. [In Thai]
- Donsakul T, Magtoon W. 2002. Karyotype of *Rasbora Caudimaculata*, *R. myersi*, *R. retrodorsalis*, *R. paviei* from Thailand. Academic conference seminar for research papers of Srinakharinwirot University. Bangkok. 1–7. [In Thai]
- Donsakul T, Magtoon W. 1995. Karyotypes of four Cyprinid fishes, *Osteochilus melanopleura*, *Puntioplites proctozyron*, *Paralaubuca riveroi* and *Rasbora sumatrana* from Thailand. In: 33rd Conference of Kasetsart University. Fisheries. 128–138. [In Thai]
- Endo A, Ingalls TH. 1968. Chromosomes of the Zebra Fish: a model for cytogenetic, embryologic, and ecologic study. *Journal Heredity* 59: 382–384. <https://doi.org/10.1093/oxfordjournals.jhered.a107755>
- Eschmeyer WN, Fricke R (Eds.) 2009. Catalog of Fishes, electronic version (updated 13 March 2009). Eschmeyer WN (Ed.) 1998. Catalog of Fishes. California Academy of Sciences, San Francisco. Available from: <<http://research.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>>.
- Filcek KB, Gilmore SA, Scribner KT, Jones ML. 2005. Discriminating lamprey species using multilocus microsatellite genotypes. *North American Journal of Fisheries Management* 25: 502–509. <https://doi.org/10.1577/M03-206.1>
- Fontana F, Chiarelli B, Rossi AC. 1970. II cariotipo di alcune specie di Cyprinidae, Centrarchidae, Characidae studiate mediante culture 'in vitro'. *Caryologia* 23: 549–564. <https://doi.org/10.1080/00087114.1970.10796394>
- Frame L, Dickerson RL. 2006. Fish and wildlife as sentinels of environment contamination. In *Endocrine disruption: biological bases for health effects in wildlife and humans*. Oxford University Press, New York, U.S.A.
- Froese R, Pauly D. 2012. Fish Base. World Wide Web electronic publication, www.fishbase.org, version 06/2019.
- Getlekha N, Molina WF, Cioffi MB, Yano CF, Maneechot N, Bertollo LAC, Supiwong W, Tanomtong A. 2016. Repetitive DNAs highlight the role of chromosomal fusions in the karyotype evolution of *Dascyllus* species (Pomacentridae, Perciformes). *Genetica* 144(2): 203–211. <https://doi.org/10.1007/s10709-016-9890-5>
- Gornung E, Gabrielli I, Cataudella S, Sola L. 1997. CMA3- banding pattern and fluorescence *in situ* hybridization with 18S rRNA genes in zebrafish chromosomes. *Chromosome Research* 5: 40–46. <https://doi.org/10.1023/A:1018441402370>
- Goldstein DB, Pollock DD. 1997. Lauching microsatellites. *Journal of Heredity* 88: 335–342. doi: 10.1093/oxfordjournals.jhered.a023114
- Jurka J, Kapitonov VV, Pavlicek A, Klonowski P, Kohany O, Walichiewicz J. 2005. Repbase Update, a database of eukaryotic repetitive elements. *Cytogenetics Genome. Research* 110: 462–467. <https://doi.org/10.1159/000084979>

- Kaewtip J, Wongwut A, Iamchuen N, Pimmongkholgul S, Seetapan K. 2021. The Standard Karyotypes and Idiograms of Rose Danio (*Danio roseus*) and Laos Danio (*Devario laoensis*). *Burapa Science Journal* 26(2): 783-789. [in Thai]
- Khuda-Bukhsh AR. 1979. Karyology of two species of hill-stream fishes, *Barilius bendelisis* and *Rasbora daniconius* (Fam. Cyprinidae). *Current Science* 48: 793-794.
- Kubat Z, Hobza R, Vyskot B, Kejnovsky E. 2008. Microsatellite accumulation in the Y chromosome of *Silene latifolia*. *Genome* 51: 350-356. <https://doi.org/10.1139/G08-024>
- Levan A, Fredga K, Sandberg AA. 1964. Nomenclature for centromeric position on chromosomes. *Hereditas* 52: 201-220. <https://doi.org/10.1111/j.1601-5223.1964.tb01953.x>
- Liehr T. 2009. *Fluorescence in situ Hybridization (FISH)-Application Guide*. Springer, Verlag, Berlin, Heidelberg. <https://doi.org/10.1007/978-3-540-70581-9>
- Manna GK, Khuda-Bukhsh AR. 1977. Karyomorphology of cyprinid fishes and cytological evaluation of the family. *Nucleus* 20: 119-127.
- Maneechot N, Yano CF, Bertollo LAC, Getlekha N, Molina WF, Ditcharoen S, Tengjaroenkul B, Supiwong W, Tanomtong A, Cioffi MB. 2016. Genomic organization of repetitive DNAs highlights chromosomal evolution in the genus *Clarias* (Clariidae, Siluriformes). *Molecular Cytogenetics* 9(1): 4. <https://doi.org/10.1186/s13039-016-0215-2>
- McCusker MR, Paterson IG, Bentzen P. 2008. Microsatellite markers discriminate three species of North Atlantic wolffishes (*Anarhichas* spp.). *Journal of Fish Biology* 72: 375-385. <https://doi.org/10.1111/j.1095-8649.2007.01701.x>
- Moraes RLR, Bertollo LAC, Marinho MMF, Yano CF, Hatanaka T, Barby FF, Troy WP Cioffi MB. 2017. Evolutionary relationships and cytotaxonomy considerations in the genus *Pyrrhulina* (Characiformes, Lebiasinidae). *Zebrafish* 14(6): 536-546. <https://doi.org/10.1089/zeb.2017.1465>
- Moraes RLR, Sember A, Bertollo LAC, de Oliveira EA, Ráb P, Hatanaka T, Marinho MMF, Liehr T, Al-Rikabi ABH, Feldberg E, Viana PF, Cioffi MB. 2019. Comparative cytogenetics and neo-Y formation in small-sized fish species of the genus *Pyrrhulina* (Characiformes, Lebiasinidae). *Frontiers in Genetics* 10: e678. <https://doi.org/10.3389/fgene.2019.00678>
- Pijnacker LP, Ferwerda MA. 1995. Zebrafish chromosome banding. *Genome* 38: 1052-1055. <https://doi.org/10.1139/g95-140>
- Pinkel D, Straume T, Gray J. 1986. Cytogenetic analysis using quantitative, high sensitivity, fluorescence hybridization. *Proceedings of the National Academy of Sciences of the United States of America* 83: 2934-2938. <https://doi.org/10.1073/pnas.83.9.2934>
- Phillips RB, Reed KM. 2000. Localization of repetitive DNAs to zebrafish (*Danio rerio*) chromosomes by fluorescence *in situ* hybridization (FISH). *Chromosome. Research* 8: 27-35. <https://doi.org/10.1023/A:1009271017998>
- Phimphan S, Chaiyasan P, Suwannapoom C, Reungsing M, Juntaree S, Tanomtong A, Supiwong W. 2020. Comparative karyotype study of three Cyprinids (Cyprinidae, Cyprininae) in Thailand by classical cytogenetic and FISH techniques. *Comparative Cytogenetics* 14(4): 597-612. <https://doi.org/10.3897/CompCytogen.v14i4.54428>
- Post A. 1965. Vergleichende Untersuchungen der Chromosomenzahlen bei Süsswasserteleosteen. *Journal Zoological Systematics and Evolutionary Research* 3: 47-93. <https://doi.org/10.1111/j.1439-0469.1965.tb00426.x>
- Racey PA, Barratt EM, Burland TM, Deaville R, Gotelli D, Jones G, Piertney SB. 2007. Microsatellite DNA polymorphism confirms reproductive isolation and reveals differences in population genetic structure of cryptic pipistrelle bat species. *Biological Journal of the Linnean Society* 90: 539-550. <https://doi.org/10.1111/j.1095-8312.2007.00746.x>
- Raskovic B, Poleksic V, Zivic I, Spasic M. 2010. Histology of carp (*Cyprinus carpio*, L.) gills and pond water quality in semiintensive production. *Bulgarian Journal of Agricultural Science* 16: 253-262.
- Reddy PB, Rawat SS. 2013. Assessment of aquatic pollution using histopathology in fish. *International Journal of Environmental Science and Technology* 2: 79-82.
- Rishi KK (1976) Karyotypic studies on four species of fishes. *Nucleus* 19: 95-98.
- Rüber L, Kottelat M, Tan HH, Ng PKL, Britz R. 2007. Evolution of miniaturization and the phylogenetic position of *Paedocypris*, comprising the world's smallest vertebrate. *BMC Evolution Biology* 7: 38. <https://doi.org/10.1186/1471-2148-7-38>
- Saenjundaeng P, Cioffi MB, Oliveira EA, Tanomtong A, Supiwong W, Phimphan S, Collares-Pereira MJ, Sember A, Bertollo LAC, Liehr T, Yano CF, Hatanaka T, Ráb P. 2018. Chromosomes of Asian cyprinid fishes: cytogenetic analysis of two representatives of small paleotetraploid tribe Probarbini. *Molecular Cytogenetics* 11(1): 51. <https://doi.org/10.1186/s13039-018-0399-8>
- Sassi FMC, Oliveira EAD, Bertollo LAC, Nirchio M, Hatanaka T, Marinho MMF, Moreira-Filho O, Arou-

- tiounian R, Liehr T, Al-Rikabi ABH, Cioffi MB. 2019. Chromosomal evolution and evolutionary relationships of *Lebiasina* species (Characiformes, Lebiasinidae). *International Journal of Molecular Sciences* 20(12): 2944. <https://doi.org/10.3390/ijms20122944>.
- Schreeb KH, Groth G, Sachsse W, Freundt KJ. 1993. The karyotype of the zebrafish. *Journal of Experimental Animal Science* 36: 27–31.
- Seetapan K, Moeikum T. 2004. Karyotypes of ten Cyprinid fishes (Family Cyprinidae). *Journal of Agricultural Research and Extension* 22: 92–101. [in Thai]
- Sharma KK, Sharma OP, Tripathi NK. 1998. Female heterogamety in *Danio rerio* (Cypriniformes: Cyprinidae). *Proceedings of the National Academy of Sciences, India* 68: 123–126.
- Sola L, Gornung E. 2001. Classical and molecular cytogenetics of the zebrafish, *Danio rerio* (Cyprinidae, Cypriniformes): an overview. *Genetica* 111: 397–412. <https://doi.org/10.1023/A:1013776323077>
- Spoz A, Boron A, Porycka K, Karolewska M, Ito D, Abe S, Kirtiklis L, Juchno D. 2014. Molecular cytogenetic analysis of the crucian carp, *Carassius carassius* (Linnaeus, 1758) (Teleostei, Cyprinidae), using chromosome staining and fluorescence in situ hybridisation with rDNA probes. *Comparative cytogenetics* 8(3): 233–248. <https://doi.org/10.3897/CompCytogen.v8i3.7718>
- Sukham S, Chingakham B, Thoidingjam L, Waikhom G. 2013. Cytogenetic characterization of *Devario aequipinnatus* (McClelland, 1839) and *Devario yuensis* (Arunkumar and Tombi, 1998) (Cypriniformes: Cyprinidae) from Manipur, northeast India. *Turkish Journal of Zoology*. 37(6): 706–712. <https://doi.org/10.3906/ZOO-1211-12>
- Supiwong W, Liehr T, Cioffi MB, Chaveerach A, Kosyakov N, Pinthong K, Tanee T, Tanomtong A. 2014. Chromosomal evolution in naked catfishes (Bagridae, Siluriformes): A comparative chromosome mapping study. *Zoologischer Anzeiger* 253(4): 316–320. <https://doi.org/10.1016/j.jcz.2014.02.004>
- Supiwong W, Saenjundaeng P, Maneechot N, Choos-eangjaew S, Pinthong K, Tanomtong A. 2017. A discovery of nucleolar organizer regions (NORs) polymorphism and karyological analysis of Crystal eye catfish, *Hemibagrus wyckii* (Siluriformes, Bagridae) in Thailand. *Cytologia* 82(4): 403–411. <https://doi.org/10.1508/cytologia.82.403>
- Supiwong W, Pinthong K, Seetapan K, Saenjundaeng P, Bertollo LAC, de Oliveira EA, Yano CF, Liehr T, Phimphan S, Tanomtong A, Cioffi MB. 2019. Karyotype diversity and evolutionary trends in the Asian swamp eel *Monopterus albus* (Synbranchiformes, Synbranchidae): a case of chromosomal speciation? *BMC Evolution Biology* 19(1): 73. <https://doi.org/10.1186/s12862-019-1393-4>
- Tang KL, Agnew MK, Hirt MV, Sado T, Schneider LM, Freyhof J, Sulaiman Z, Swartz E, Vidthayanon C, Miya M, Saitoh K, Simons AM, Wood RM, Mayden RL. 2010. Systematics of the subfamily Danioninae (Teleostei: Cypriniformes: Cyprinidae). *Molecular Phylogenetics Evolution* 57(1): 189–214. <https://doi.org/10.1016/j.ympev.2010.05.021>.
- Tautz D, Renz M. 1984. Simple sequences are ubiquitous repetitive components of eukaryotic genomes. *Nucleic Acids Research* 25: 4127–4138. <https://doi.org/10.1093/nar/12.10.4127>
- Terencio ML, Schneider CH, Gross MC, Vicari MR, Farias IP, Passos KB, Feldberg E. 2013. Evolutionary dynamics of repetitive DNA in *Semaprochilodus* (Characiformes, Prochilodontidae): a fish model for sex chromosome differentiation. *Sexual Development* 7(6): 325–333. <https://doi.org/10.1159/000356691>
- Ueda T, Naoi H. 1999. BrdU-4Na-EDTA-Giemsa band karyotypes of three small freshwater fish, *Danio rerio*, *Oryzias latipes*, and *Rhodeus ocellatus*. *Genome* 42: 531–535. <https://doi.org/10.1139/g98-153>
- Vidthayanon C. 2005. Thailand Red Data: Fishes. Office of Natural Resources and Environmental Policy and Planning. Bangkok. [in Thai]
- Yeesaem N, Jantararat S, Yeesin P. 2019. Cytogenetic Characterization of *Rasbora einthovenii* in Sirindhorn Peat Swamp Forest, Narathiwat Province. *Journal Fish Technology Research* 13: 58–68. [in Thai]
- Yano CF, Poltronieri J, Bertollo LAC, Artoni RF, Liehr T, Cioffi MB. 2014. Chromosomal mapping of repetitive DNAs in *Triplotheus trifurcatus* (Characidae, Characiformes): insights into the differentiation of the Z and W chromosomes. *PLoS ONE* 9(6): e90946. <https://doi.org/10.1371/journal.pone.0090946>
- Yano CF, Bertollo LAC, Cioffi MB. 2017. Fish-FISH: Molecular cytogenetics in fish species. In: Liehr T (ed.), *Fluorescence in situ Hybridization (FISH)-Application Guide*, 429–443. Springer, Berlin, Germany.
- Yenchum W. 2010. Histological effects of carbofuran on guppy *Poecilia reticulata* Peters. Ph.D Thesis in Biological Science, Chulalongkorn University, Bangkok, Thailand. [in Thai]